



WATER QUALITY

NEW ZEALAND PLANTED FORESTS ENVIRONMENTAL FACTS

There are some 1.7 million hectares of planted forests in New Zealand, 90% of which are in radiata pine.

These forests are home to an estimated 24,220 kms of streams that, for most of the forest growing cycle, provide a source of high quality water to downstream users.

There is increasing pressure on our water resources, however with prudent stewardship and ongoing improvements to management practices, New Zealand's forests will continue to provide sustainable sources of high quality water.

FORESTS GENERATE HIGH QUALITY WATER

Water quality is a term used to describe the physical, chemical and biological characteristics of water, and is influenced by the long term cyclic nature of planted forests in New Zealand.

Forests have a strong influence on the water quality of catchments. They moderate the local climate, which in turn influences the quantity, temperature and overall quality of stream water. As a result, planted forests almost always yield better water quality than other developed land uses, including agriculture and urban development.

Afforestation. Planting forests can improve the water quality from land that was previously in pasture, within 5-6 years, by reducing water temperature, nutrient levels and sediment yields. This highlights the potential of forestry as a remedial tool for degraded waterways.

Over a rotation. The largest part of the forest cycle is in mid- to mature rotation. Water quality during this phase is naturally variable, but is characterised by cool water temperature, low sediment and nutrient concentrations, and healthy invertebrate communities.

Mature planted forests. Have similar water quality attributes to those found in undisturbed indigenous forests. Streams are often cool and shaded, and are strongly influenced by the riparian vegetation type and channel width. The low light and cooler temperatures promote high levels of dissolved oxygen in the water, which is a good indicator of water quality and essential to supporting the aquatic environment.

Actively managing harvesting. Harvesting can adversely affect water quality through increased sunlight and temperature. However, residual shade provided by the stream bank, logging slash, surrounding topography and any remaining vegetation all have an effect on maintaining water quality attributes, with shade recovery dependent on the size of the stream and the regrowth of riparian vegetation.

Harvesting in steep-land country can increase sediment yields reaching waterways from roading and post-harvest erosion. Sediment yields elevated by harvesting activities typically return to pre-harvest levels, or decline markedly, within two to six years of harvest.

Harvesting impacts can be modified by retaining riparian management zones and to a lesser extent, retaining moderate quantities of logging slash across small stream channels. In addition, minimising tracking, good hauler set-up and implementation, and applying sediment control structures and techniques to the earthworks associated with harvesting.

Retaining intact riparian management zones during harvest helps maintain low light levels and temperatures. Riparian zones can reduce the amount of harvesting debris deposited in the stream and contribute to bank stability. They are limited in their ability to filter out point source sediment and any associated sediment-bound phosphorus. Riparian management zones can also limit nitrate levels following harvest, helping to maintain the aquatic community at similar levels to those found in mature pine and indigenous forest streams.

THE CHALLENGES OF STEEPLAND FORESTRY

Around a third of New Zealand's planted forest is in steep hill country (>25°), which is often highly erodible and susceptible to extreme weather events. Steepland forests have a higher density of stream networks than those on flatter topography. These networks extend into the upper reaches of headwater catchments, and when subjected to intense rainfall, can result in flooding and debris flows.

Steepland harvesting. Reducing the total length of the roading infrastructure will reduce the potential for sediment generation, however there is a trade-off between roading length, keeping riparian buffers intact and logging across the stream channel.

Reduced road length and harvesting across stream channels can provide better water quality outcomes, although there will still be some impact on water quality, habitat and a range of other biological processes. It is preferable to prevent the export of sediment directly or indirectly into a river.

Recovery time depends on site conditions, harvest method, post-harvest management techniques and the scale of operations.

Post-harvest activities. A number of site establishment and silvicultural activities, such as pesticide (including herbicide) and fertiliser use, may affect water quality. Compared to other productive land uses, these activities are used infrequently during the growing phase of a production forest crop, and when applied using good practice, the risk of impacting water quality is low.

Aerial application in rugged terrain with dense stream networks can increase the risk of impacting water quality.



MANAGING FOR MAXIMUM BENEFIT

Planted forests produce high water quality for a large component of the forestry cycle, providing a valuable community service to downstream users. The period during harvest requires the most sensitive management.

To realise the maximum benefit, the forest sector operates under the RMA (Resource Management Act) plus a range of sector-developed manuals; the New Zealand Roding Manual; New Zealand Environmental Code of Practice; individual company environmental management systems for good practice for maximum environmental outcomes; and, in some cases, Forest Stewardship Council (FSC) standards.

CONTACT
Email: publications@scionresearch.com Phone: +64 7 343 5899

KEY LINKS
www.youtube.com/user/ScionResearch 'Forests and Water: Creating prosperity from trees'.
Growing confidence in forestry's future: www.gcff.co.nz
Ministry for the Environment overview of fresh water: www.mfe.govt.nz/fresh-water/overview-fresh-water/about-fresh-water-nz

KEY REFERENCES
Baillie, BR, & Neary, DG (2015). Water quality in New Zealand's planted forests: a review. *New Zealand Journal of Forestry Science*, 45(1).
Baillie, BR, Rolando, CA. (2015). Long-term management of streams in planted forest steep-lands. *New Zealand Journal of Forestry*, 60(2), 21-24.
New Zealand Forest Owners Association (2007). *New Zealand Environmental Code of Practice for Plantation Forestry* (Version 1).
New Zealand Forest Owners Association (2011). *New Zealand Forest Road Engineering Manual*.